

# Effect of Some Fermentation Substrates and Growth Temperature on Exopolysaccharide Production by *Streptococcus Thermophilus* BN1

Bennama Rabha, Rechidi-Sidhoum Nadra, and Bensoltane Ahmed

**Abstract**—The purpose of this research was to evaluate, the effect of incubation temperature and some substrates on exopolysaccharides (EPS) production. The *S. thermophilus* BN1 an EPS-producer strain isolated from cow milk was examined in this study. Skimmed milk, whole milk and cheese whey, were selected as culture media and two different temperatures were also tested. The EPS produced in the different studied conditions was purified and quantified. The strain BN1 was able to produce EPS in all established conditions. However, significant differences were observed in the amounts of produced EPS. The strain has been shown to produce high amounts of EPS at 37°C compared to 42°C ( $P < 0.05$ ). In skimmed milk at 37°C, the EPS amounts reached 548 PDM mg l-1, followed by 375 PDM mg l-1 and 325 PDM mg l-1 respectively in whey and whole milk. In addition, a slight significant difference was noted on the biomass, pH and lactic acid values obtained from the three fermented substrates ( $p < 0.05$ ). The present results demonstrate that suboptimal growth temperature (37°C) had a significant effect on the EPS production by *S. thermophilus* BN1.

**Index Terms**—*S. thermophilus*, exopolysaccharide production, growth temperature, substrate.

## I. INTRODUCTION

Today, microbial polysaccharides have regarded as interesting natural bio-ingredients in many industrial sectors. Their biological and rheological properties are widely used particularly in food sector, where they are used as gelling, emulsifiers and stabilizers agents [1]. Lactic acid bacteria (LAB) play a key role during the fermentation process since they contribute to the texture, flavour, quality and conservation of the fermented products. Several strains of LAB, are also able to produce exopolysaccharides (EPS), the main advantage of EPS from LAB is that they are produced by food-grade microorganisms known as GRAS (Generally Recognized as Safe) compared to other microbial EPS [2, 3]. These compounds have attracted great interest since they can act as natural thickeners that improve the texture properties, decrease syneresis and reduce the fat levels in fermented dairy foods. According to their, chemical composition the EPS can be classified into two major types: the homopolysaccharides and heteropolysaccharides [2]. Some

EPS may remain attached to the cell in capsule form [4]. Bacterial capsules are viscous and can affect the physical properties of cultured milk [5]. EPS-producing LAB belong to different genera such as *Bifidobacterium*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Pediococcus* and *Streptococcus* [6]. The observations made to date on the conditions for EPS production by LAB do not appear to be generalized, but rather related to a species, or even each studied strain [7].

Indeed, optimization designed to determine the operating growth conditions that tend to improve performance of interesting bacteria. In this work we focus on the production of EPS by *S. thermophilus* BN1. *S. thermophilus* is one of the main starters used in the dairy industry for the manufacture of cheese and yogurt, came in the second place after *Lc. lactis* [3]. Its EPS production has been related to the growth phase and the culture conditions [8]. Whereas in fermented milks, EPS concentrations were less than 600 mg l<sup>-1</sup>, M17 media yielded up to 1500 mg l<sup>-1</sup> depending on the nature of the carbohydrate source used and the carbon/nitrogen ratio [9, 10]. These data show the importance of growth conditions on EPS production.

Regarding the topic of EPS production, in this paper, the influence of the growth temperature and the type of fermentation media on EPS production by *S. thermophilus* BN1 will be analyzed.

## II. MATERIAL AND METHODS

### A. Strain, Media and Growth Conditions

*S. thermophilus* BN1, a strain isolated from raw milk [11] was grown in M17 supplemented with lactose (2% w/v) (LM17) at 42°C under anaerobic conditions. EPS production by this strain was evaluated using as substrates: skimmed milk, whole milk, (Candia, Algeria) and cheese whey collected from the cheese factory “Sidi Saâda” (Relizane, Algeria). The preparation of the whey was performed using the protocol described by Shams and Jaynes [12]. The residues of milk proteins present in the whey samples have been partially precipitated and the medium was adjusted at the same pH of milk substrates. All substrates were heated for 5min. at 95°C and then cooled to the temperature of incubation.

### B. Growth on Different Fermentation Substrates and Eps Determination

EPS production was determined in cultures performed in skimmed milk, whole milk and cheese whey. 250 ml of each

Manuscript received December 30, 2011; revised January 30, 2012.

Bennama Rabha, Rechidi-Sidhoum Nadra are with the Département de Biotechnologie, Université de Mostaganem, BP 227, 27000. Mostaganem, Algeria (Tel.: 00 213 771623291; e-mail: bennamarabha@yahoo.fr).

Bensoltane Ahmed is with Département de Biologie, Université d'Oran, Algeria.

medium was inoculated at 0.5% (v/v) with *S. thermophilus* BN1 previously grown in LM17 as has been described before. These cultures were incubated at 37°C and 42°C for 17 h and 12h respectively. Then, aliquots were taken from each medium to determine different parameters mentioned below.

### C. Growth Evaluation

From the established cultures, serial dilutions were made in peptone saline water [(1 g.l<sup>-1</sup>) and NaCl (8.5 g.l<sup>-1</sup>)]. Appropriate dilutions were spread on LM17 agar and incubated for 48 h at 42°C. Growth was estimated as log CFU/ml.

### D. Measurement of Total Titratable Acidity (TTA) and PH

The concentration of lactic acid in the media was determined by titration with N/9 NaOH solution. The pH was also measured with a pH meter type "Hanna instruments microprocessor pH meter".

### E. Isolation and Purification of EPS

Isolation of EPS was performed using the protocol described by Salazar *et al.* [13]. Briefly, this protocol is based on proteins elimination by precipitation with TCA (12 %) and subsequently precipitation of EPS using two volumes of cold ethanol. The Precipitated EPS crude was recuperated after centrifugation (10000xg, 4°C, 30 min.) These samples were then purified by dialysis against water using a membrane (Sigma) with a molecular cut-off weight of 12-14 kDa for three days at 4°C.

### F. Quantification of the Purified EPS

EPS quantification was done by gravimetric analysis [8], the polymer dry mass (PDM) of the purified EPS; was determined after 48h of drying at 42°C. The measured values are subtracted from that obtained with a PDM of the control media to get real amounts of EPS produced by the strain. Also, the presence of capsule like-EPS was visualised by negative staining technique using India ink.

### G. Data Analysis

All experiments were repeated two times. The results are means of two replicates. The Student's t-test was applied to compare the means values of the measured parameters.

## III. RESULTS AND DISCUSSION

### A. Bacterial Growth and Acidification Activity

Cultured on the three tested substrates, *S. thermophilus* BN1 was able to grow and show relatively high number of viable cells at the end of each fermentation, with an average increase of 2.5 log CFU/ml compared to the initial rate of inoculation. The biomass developed after 12h of incubation at 42°C and 17h at 37°C is shown in Fig 1. At 37°C in skimmed milk, the biomass was quite similar to that obtained in whole milk in the conditions analyzed. By contrary, at 42°C in whey it was slightly higher to that posted at 37°C. It is also interesting to note that in whole milk the difference between the growth at 42°C is bigger than in whey or skimmed milk (P<0.05). The increase in biomass caused a significant acidification in the three studied media. This was

confirmed by TTA and pH values measured after 12h and 17h of fermentation (Fig.2, 3). The acidification in the cultures of skimmed and whole milks provoked the clotting of the milk. However, in the whey culture a thick aspect was observed. The ability of *S. thermophilus* BN1 to grow on the tested media is, in fact, due to the capacity of this species to metabolize lactose present naturally in these media. Its conversion into lactic acid decreases the final pH of the medium [14]. By HPLC analysis of some fermented skimmed milk samples, it was found that *S. thermophilus* BN1 also produced small amounts of other acids such as formic and acetic acids (data not shown).

### B. Effect of Incubation Temperature on Growth and Eps Production

The growth temperature is considered, the most important factor influencing and having a significant impact on EPS production in *S. thermophilus* [15]. In this context, the effect of temperature on EPS production was analyzed. EPS production was observed at both temperatures. The higher EPS production was reached at 37°C compared to 42°C independently of the media culture. At 37°C amounts of 548 mg PDM l<sup>-1</sup>, 375 mg PDM l<sup>-1</sup> and 325 mg PDM l<sup>-1</sup> were produced in skimmed milk, whey and whole milk respectively (Fig. 4). These observed results reveal the existence of significant effect of incubation temperature (37°C) on the EPS biosynthesis by *S. thermophilus* BN1. However, a well-documented finding showed that 37°C is a suboptimal growth temperature of *S. thermophilus*. Several studies have been confirmed, that suboptimal temperatures influence positively EPS production by mesophilic and thermophilic LAB [6, 16]. Nevertheless, de Vuyst *et al.* [8] obtained maximal EPS production, when *S. thermophilus* grew at its optimum temperature (42°C). These data show that EPS production could vary among the different strains of *S. thermophilus* under similar growth conditions.

This characteristic is not specific for *S. thermophilus*, a large diversity in EPS production was also observed in other LAB for instance in *Lactobacillus delbrueckii ssp. bulgaricus* from 57 to 424 mg l<sup>-1</sup> [17], 30-85 mg l<sup>-1</sup>, 100-600 mg l<sup>-1</sup> and 105-150 mg l<sup>-1</sup> in *L. lactis subsp. lactis* or, *L. lactis subsp. cremoris* and *Lb. casei subsp. casei*, respectively [16] and 105-168 mg l<sup>-1</sup> with bifidobacteria [18]. It is also interesting to note that at 37°C the fermented skimmed milk had a firm and compact texture difficult to break. This textural aspect was reported by many authors in other EPS-producing LAB strains incubated at suboptimal temperatures. The EPS overproduction at suboptimal temperature has been proposed as answer mechanism to the physiological stress at these temperatures, especially on that species such as *S. thermophilus* with a deficient proteolytic system. In addition, Vaningelgem *et al.* [19] reported that the use of high-level EPS-producing strain displaying a stronger milk-clotting ability. However, in thermophilic LAB, the amount of EPS produced is often related to growth, therefore, to the produced biomass [8].

### C. Effect of Fermentation Media on Growth and Eps Production

Several authors have stated that medium composition either carbon source, nitrogen source, or ion source are important parameters in EPS biosynthesis [6, 19]. The results

have showed that BN1 strain is able to grow and produce EPS in all substrates (Fig 4). The maximal EPS production occurred at 37°C in skimmed milk with 548 PDM mg l<sup>-1</sup> compared to whey (375 PDM mg l<sup>-1</sup>) and whole milk (325 PDM mg l<sup>-1</sup>). The obtained results are according to the findings of de Vuyst *et al.* [8] indicate that milk is the best media for EPS production in *S. thermophilus* compared with other media (MRS, M17 or SDM). Other authors [19] have observed that EPS production in some strains of *S. thermophilus* increases only when the skim-based media are enriched with peptone or yeast extract. Moreover, at 42°C the EPS production by *S. thermophilus* BN1 in the used substrates was approximately three times lower to that observed at 37°C (Fig. 4).

In this study, the interesting result was the ability of *S. thermophilus* BN1 to produce EPS in whey medium. Most studies [20] on EPS production by *S. thermophilus* were realized using whey, but only as based medium and not as a single substrate. Our results demonstrate the capacity of BN1 strain to produce EPS using whey as a substrate without exogenous nutrients. Whey is considered to be a rich medium; besides lactose it provides amino acids. The presence of amino acids is essential to EPS production by *S. thermophilus* BN1, the enrichment of milk with amino acids yielded a significant increase on EPS production (data not shown). According to the literature, the amino acids can generate the metabolic energy, based on the production of ATP through the phosphorylation and decarboxylation of the substrates and the transport of generator, respectively [21]. This energy is essential for cell growth and growth-associated with EPS production. The production of EPS in whey has a great economic interest because; it allows to upgrade this product that is usually rejected by most of dairy industries in Algeria. On the other hand, in the three fermented media a shooting or slime texture was observed at 42°C. Moreover, this observed aspect of fermented media obtained under this temperature could reveal a homogeneous EPS production. While at 37°C, a firm texture has characterized the samples of fermented skimmed milk. These textural characteristics could be related to the production of two types of EPS, previously observed in this strain [11]. Ever and about the texture, the microscopic observations performed on samples taken from fermented substrates showed the presence of capsule in this strain (Fig. 5). In fact, the capsule formation in this strain was already described by Bennama *et al.* [11]. The influence of this type of capsular EPS on the texture of fermented dairy products was confirmed by many authors [4, 22]. For this purpose, the presence of capsule-like EPS in this strain could also explain the different textures observed on the fermented substrates.

However, Lemoine *et al.* [23] suggested that the textural properties of several EPS-producing *S. thermophilus* are due to the synthesis of polysaccharides of similar chemical composition, but different structure. In dairy industry, the firm and shooting textures are very important. At this scale, an appropriate processing allows the interaction of the EPS with milk proteins, somehow breaking the protein matrix. Once broken, it would modulate the organoleptic properties of the end-product.

#### IV. CONCLUSION

From all these considerations, it appears that BN1 strain produces secreted and capsular EPS in all the tested substrates. Also, this study has demonstrated that the EPS production by *S. thermophilus* BN1 is depending on the growth conditions; i.e. the incubation temperature (37°C), which stimulated strongly the EPS production. Similarly, the nature of the fermentation substrate has also influence on EPS production. These factors associated may give optimal growth rate and have positive effects on the organoleptic quality of the fermented product. EPS production by *S. thermophilus* BN1 strain in whey substrate shows promising possibilities to promote this waste industrial dairy product.




Fig. 1. Biomass developed (log CFU/ml) by BN1 strain after 12h and 17h of incubation in the fermented substrates. SM: Skimmed milk; WM: Whole milk; Wh: whey; CS: Control Skimmed milk; CWM: Control whole milk; CWh: Control whey.




Fig. 2. PH values measured after 12h and 17h of incubation in the substrates fermented by BN1 strain. SM: Skimmed milk; WM: Whole milk; Wh: whey; CS: Control Skimmed milk; CWM: Control whole milk; CWh: Control whey




Fig. 3. Lactic acid (mmol/l) produced by BN1 strain after 12h and 17h of incubation in the fermented substrates. SM: Skimmed milk; WM: Whole milk; Wh: whey; CS: Control Skimmed milk; CWM: Control whole milk; CWh: Control whey.




Fig. 4. EPS amounts (mg PDM l<sup>-1</sup>) produced by BN1 strain after 12 h and 17h of incubation in the fermented substrates. SM: Skimmed milk; WM: Whole milk; Wh: whey




Fig. 5. Cells of BN1 strain surrounded by clear halo showing the formation of capsule -like EPS - Negative contrast obtained by India ink.

#### REFERENCES

- [1] A. Laws, Y. Gu, and V. Marshall, "Biosynthesis, characterisation and design of bacterial exopolysaccharides from lactic acid bacteria," *Biotechnol Adv.*, vol. 19, 2001, pp. 597-625.
- [2] B. Degeest, and L. De Vuyst, "Heteropolysaccharides from lactic acid bacteria," *FEMS Microbiol Review.*, vol. 23, 1999, pp. 153-177.
- [3] C. Delorme, "Safety assessment of dairy microorganisms: *Streptococcus thermophilus*," *Int J Food Microbiol.*, vol. 126, 2008, pp 274- 277.
- [4] A. N. Hassan, "Possibilities and Challenges of Exopolysaccharide-Producing Lactic Cultures in Dairy Foods," *J Dairy Sci.*, vol. 91, 2008, pp1282-1298.
- [5] A. N. Hassan, J. F. Frank, K. A. Schmidt, and S. Shalabi, "Rheological Properties of Yogurt Made with Encapsulated Nonropy Lactic Cultures," *J Dairy Sci.*, vol. 79, 1996, pp. 2091-2097.
- [6] J. Cerning, "Exocellulair polysaccharides produced by lactic acid bacteria," *FEMS Microbiol Reviews.*, vol. 87, 1990, pp. 113-130.
- [7] G. Corrieu, and F. M. Luquet, "Bactéries lactiques: de la génétique aux ferments," Eds. Tec and Doc. Paris (France). 2008, pp. 319-331.
- [8] L. de Vuyst, F. Vanderveken, S. Van de Ven, and B. Degeest, "Production by and isolation of exopolysaccharides from *Streptococcus thermophilus* grown in a milk medium and evidence for their growth-associated biosynthesis," *J Appl Microbiol.*, vol. 84, 1998, pp. 1059-1068.
- [9] J. Cerning, C. M. G. C. Renard, J. F. Thibault, C. Bouillanne, M. Landon, M. Desmazeaud, and L. Topisirovic, "Carbon source requirements for exopolysaccharide production by *Lactobacillus casei* CG11 and partial structure analysis of the polymer," *Appl Environ Microbiol.*, vol. 60, 1994, pp. 3914-3919.
- [10] S. Petry, S. Furlan, M. J. Crepeau, J. Cerning, and M. Desmazeaud, "Factors affecting exocellular polysaccharide production by *Lactobacillus delbrueckii* subsp. *bulgaricus* grown in a chemically defined medium," *Appl Environ Microbiol.*, vol. 66, 2000, pp. 3427-3431.
- [11] R. Bennama, M. Fernández, V. Ladero, M. A. Alvarez, N. Rechidi-Sidhoum, and A. Bensoltane, "Isolation of an exopolysaccharide-producing *Streptococcus thermophilus* from Algerian raw cow milk," *Eur Food Res Technol.*, vol. 234, 2012, pp. 119-125.
- [12] M. A. Shams, and H.O. Jaynes, "Characterization of exopolysaccharides produced by *Corynebacterium* #98 in cheese whey substrate," *J Food Sci.*, vol. 48, 1983, pp. 208-211.
- [13] N. Salazar, A. Prieto, J. A. Leal, B. Mayo, J. C. Bada-Gancedo, C. G. de los reyes-Gavilán, and P. Ruas-Madiedo, "Production of exopolysaccharides by *Lactobacillus* and *Bifidobacterium* strains of human origin, and metabolic activity of the producing bacteria in milk," *J Dairy Sci.*, vol. 92, 2009, pp. 4158-4168.
- [14] A. Zourari, S. Roger, C. Chabanet, and M. J. Desmazeaud, "Characterisation of lactic acid bacteria from Greek yaourt," *Lait.*, vol.71, 1991, pp. 445-461.
- [15] A. S. Kumar, K. Mody, and B. Jha, "Bacterial exopolysaccharides- a perception," *J Basic Microbiol.*, vol. 47, 2007, pp. 103-117.
- [16] J. Cerning, C. Bouillanne, M. Landon, and M. Desmazeaud, "Isolation and characterization of exopolysaccharides from slime-forming mesophilic lactic acid bacteria," *J Dairy Sci.*, vol. 75, 1992, pp. 692-699.
- [17] J. Cerning, C. Bouillanne, M. J. Desmazeaud, and M. Landon, "Isolation and characterization of exocellular polysaccharide produced by *Lactobacillus bulgaricus*," *Biotechnol Letters.*, vol. 8, 1986, pp. 625-628.
- [18] C. M. Roberts, W. F. Fett, S. F. Osman, C. Wijey, J. V. O'Connor, and D. G. Hoover, "Exopolysaccharide production by *Bifidobacterium longum* BB-79," *J Appl Bacteriol.*, vol. 78, 1995, pp. 463-468.
- [19] F. Vaningelgem M. Zamfir, F. Mozzi, T. Adriany, M. Vancanneyt, J. Swings, and L. De Vuyst, "Biodiversity of Exopolysaccharides produced by *Streptococcus thermophilus* strains is reflected in their production and their molecular and functional characteristics," *Appl Environ Microbiol.*, vol. 70, 2004, pp. 900- 912.
- [20] A. Ricciardi, E. Parente, M. A. Crudele, F. Zanetti, G. Scolari, and I. Mannazu, "Exopolysaccharide by *Streptococcus thermophilus* SY: production and preliminary characterization of the polymer," *J Appl Microbiol.*, Vol. 92, 2002, pp. 297-306.
- [21] K. Abe, H. Hayashi, and P. C. Maloney, "Exchange of aspartate and alanine mechanism for development of a proton-motive force in bacteria," *J Biol Chem.*, vol. 271, 1996, pp. 3079-3084.
- [22] A. N. Hassan, M. Corredig, and J. F. Frank, "Capsule Formation by Nonropy Starter Cultures Affects the Viscoelastic Properties of Yogurt during Structure Formation," *J Dairy Sci.*, vol. 85, 2002, pp.716-720.
- [23] J. Lemoine, F. Chirat, J. M. Wieruszkeski, G. Strecker, N. Favre, J. R. Nesser, "Structural characterization of exocellular polysaccharides produced by *Streptococcus thermophilus* SFi39 and SFi12," *Appl Environ Microbiol.*, vol. 63 (9), 1997, pp. 3512-3518.